

Indigenous vegetables consumed as lalapan by Sundanese ethnic group in West Java, Indonesia: Potential, traditions, local knowledge, and it's future

by Diana Hernawati

Submission date: 16-May-2023 11:25AM (UTC+0700)

Submission ID: 2094347342

File name: Form_B_-_1.C.1.c.pdf (2.22M)

Word count: 10584

Character count: 55278



Contents lists available at ScienceDirect

South African Journal of Botany

journal homepage: www.elsevier.com/locate/sajb



9 Indigenous vegetables consumed as *lalapan* by Sundanese ethnic group in West Java, Indonesia: Potential, traditions, local knowledge, and its future

Diana Hernawati*, Rinaldi Rizal Putra, Vita Meylani

19
Department of Biology Education, Faculty of Teacher Training and Science Education, Universitas Siliwangi, Jl. Siliwangi 24 Tasikmalaya City, 46115 Indonesia



ARTICLE INFO

24
Article History:
Received 18 August 2021
Revised 7 June 2022
Accepted 4 September 2022
Available online xxx

Edited by: A. Ndhala

Keywords:

Lalapan
Sundanese ethnic
Potential
Traditional
Local knowledge
Future

ABSTRACT

The local knowledge of indigenous people about traditional vegetables has been well documented. However, there is little conventional plant documentation biodiversity on indigenous vegetables used as lalapan in West Java, Indonesia. The method used in this study was a qualitative method using an exploratory approach. Informants, including traditional leaders, indigenous people, and the Sundanese younger generation, provided data. Credibility, transferability, dependability, and confirmability tests were used to determine the validity. The data were evaluated using data reduction, visualization, and verification techniques. Then, to show the local importance of each species, Relative Frequency of Citation (RFC) analysis was conducted, and The usage value (UV) analysis was used to determine the relative usefulness of plants in a given location. There were 86 species of edible indigenous vegetables consumed as lalapan, which belong to 32 families. The vegetables were accumulated from four traditional villages. The implication of indigenous vegetables as lalapan is a top priority to ensure nutrition, public health, and food security.

© 2022 SAAB. Published by Elsevier B.V. All rights reserved.

1. Introduction

Lalapan is a term used for indigenous vegetables consumed raw (Hendariningrum, 2018; Purba et al., 2013), especially by the Sundanese ethnic group of West Java. Edible plants that become lalapan or "fresh vegetables" may be certain wild plants eaten by humans or deliberately taken to become cultivated plants (Guerrero, 2015). Lalapan is usually served with rice and other side dishes. Lalapan is similar to salads, and some varieties are raw or fresh; others must be cooked before consumption. The Sundanese people's everyday diet is nearly inseparable from lalapan (Hendariningrum, 2018). Even when the main dish is not available, lalapan is used to substitute for the main dish by adding some sauce. Such presentation of food has become a tradition and habit of the Sundanese people, especially Sundanese people who live in traditional villages, which are rural areas whose people still maintain and uphold the customs of their ancestors.

Various kinds of plants used by the Sundanese people as salad plants are scattered in every area, especially West Java which the Sundanese people dominate. As researched by Cahyanto et al. (2018), 50 species of plants are used by the Sundanese people in the Subang Regency, grouped into 19 tribes. Then, the results of Cita's research

(2020) show that people in Nyangkewok Village, Sukabumi Regency, consume various types of lalap plants consisting of 101 plant species, most of which are dominated by plants with the Cucurbitaceae tribe. Based on the types of plants used in the two areas, there are some similarities to several kinds of plants used in the two regions, namely the number of 27 types of vegetables that are the same between the results of the research by Cahyanto et al. (2018) and Cita (2020).

Then, similar research in addition to those carried out in the two regions has also been carried out in Kampung Gunungwang and Kampung Leuweung Kolot, Bogor Regency (Amrinanto et al., 2019). The results of the research by Amrinanto et al. (2019) explained that in the two villages, there are 21 types of plants about the number of plants used as salads. The most consumed vegetables were: Cucumbers (*Cucumis sativus*); Long beans (*Vigna unguiculata*); Cassava (*Manihot esculenta*); Cabbage (*Brassica oleracea*); Poh-pohan (*Pilea melastomoides*); and Petai (*Parkia speciosa*). Consumption of fresh vegetables in these two villages is quite intense, carried out by both children and adults. Therefore, the examination results of the respondent food samples showed the high content of carotene in the blood, which is very important to support and protect human health; this is one of the factors driving the consumption of fresh vegetables among children and adults.

The Sundanese people's consumption of fresh vegetables is a positive habit in maintaining health through a healthy diet. The fitness supported by consuming fresh vegetables cannot be separated from

64
* Corresponding author.
E-mail address: hernawati@unsil.ac.id (D. Hernawati).

the active ingredients of these vegetables. The content of active ingredients in general in the form of flavonoids, terpenoids, antioxidants, and masterols have been identified in petai (*P. speciosa*) (Buanasari et al., 2017; Chhikara et al., 2018; Saleh et al., 2021; Singhania et al., 2021) and cucumber (*C. sativus*) (Uthpala et al., 2020; Uzuozokaro et al., 2018) which have anti-inflammatory, anti-hyperglycemic, and anti-microbial properties. Then, similar compounds were also found in cabbage (*B. oleracea*) (Cartea et al., 2011) and long beans (*V. unguiculata*) in addition to -sitosterol, ethanolamine, phytol, and phytosphingosine (Perchuk et al., 2020). In other indigenous communities, such as Pakistan, *Paonia emodi* has antioxidants (Ahmad et al., 2018b).

In Indonesia, there are traditional ethnic groups that have clearly-defined cultural characteristics and identities, so, likely, people's perceptions and conceptions of natural resources in their environment are different (Dewi et al., 2017). The cultural diversity of the Indonesian people is evident in kampung or villages, as well as within ethnic groups. Each region has an identity that characterizes its uniqueness (Hidayat et al., 2010). East Priangan, which is part of the West Java region, has traditional villages well known by the public and have been recognized by the government. A traditional village is a village that has ethnic characteristics with its own local culture and customs. The traditional villages in East Priangan include Naga Village in Tasikmalaya Regency (Harashani, 2018; Sudjatnika, 2018), Kuta Village in Ciamis Regency (Illiyan, 2018), and Pulo Village and Dukuh Village that are both in Garut Regency (Hidayat et al., 2010). The traditional villages in the East Priangan area are well-known for their local communities, still strong in their time-honored traditions and local knowledge (Harashani, 2018). West Java Province itself is also known as 'Tatar Sunda'. 'Tatar' has the meaning of a region with a strong culture that is still adhered to by its people.

The existence of a traditional village is always synonymous with the ties between ancestral origins, ceremonies, arts, and its organizational system, making traditional villages in each region different (Hidayat et al., 2010; Qodariah and Amiyati, 2013). The difference includes the utilization and relationships between plants, animals, and nature as part of their life. The traditional villages in East Priangan are Sundanese local identity, thus making these villages different from traditional villages in other regions. One of the distinguishing elements lies in natural resources and human resources (Alves, 2016; Hadi and Takaoka, 2018). Geographically, Tatar Sunda, located in the West Java region, is surrounded by mountains and mountains that make the land of Sunda so fertile and overgrown with various types of plants (Hendarinigrum, 2018).

The biodiversity in Tatar Sunda is a local resource for the surrounding community. The reciprocal interaction between humans and nature or their local ecosystem makes humans indigenous people (Harashani, 2018; King et al., 1996), humans who utilize and manage indigenous natural resources. Human action in treating nature depends on various individuals or communities understanding or perceiving nature. In general, indigenous people or traditional communities utilize and manage natural resources always based on a local knowledge system, commonly known as indigenous knowledge (Hidayat et al., 2010; Sujarwo and Caneva, 2016b). The local genius of local people with a belief system called 'worldview' can create an adaptive management system to utilize natural resources sustainably (Padhy et al., 2017; Wilson et al., 2017).

Therefore, there is a great need to integrate the people's knowledge in traditional villages with indigenous expertise from generation to generation. It remains a common challenge when indigenous knowledge can advance and improve the quality of human life and the quality of the surrounding environment (Sujarwo et al., 2016a). For this reason, it is considered essential to develop adaptive skills and the ability to apply knowledge and skills flexibly and creatively in different situations. It is hoped that the indigenous knowledge of local communities, especially indigenous people, can be followed up

through social interaction, reflection, and learning (Mokgaya, Mushaphi, and Tshisikhawe, 2019; Negi et al., 2017).

Several countries, such as Japan, Pakistan, Kenya, China, India, Italy, Thailand, and Turkey, have documented indigenous vegetables (medicinal plants, edible wild plants, and green vegetables) (Abbasi et al., 2013; Tufts et al., 2015; Uchiyama et al., 2017; Dalar et al., 2016; Bhatia et al., 2018; Guarrera and Savo, 2016; Kang et al., 2014; Birjees et al., 2020; Nazish et al., 2020; Majeed et al., 2021). Many studies have shown that ethnic sources can diversify diets and improve health attributes (Liu et al., 2016; Saba et al., 2018; Rehman et al., 2014). However, there is still little documentation of the biodiversity of traditional plants, especially indigenous vegetables, which are used as lalapan in Tatar Sunda, Indonesia. The implications of indigenous vegetables as lalapan as a top priority to ensure nutrition, public health, and food security are still not widely documented. For this reason, it is essential to provide information about indigenous vegetables consumed as lalapan as material to study the local potential of the local knowledge in traditional villages and about their future.

2. Methods

The study's data gathering methodologies include data validation, observation, and interviews (Creswell and Creswell, 2018; Sugiyono, 2017). Credibility, transferability, dependability, confirmability tests are all included in the data validity test (Lincoln and Guba, 1985; Thomas and Magilvy, 2011). Triangulation is used to determine the authenticity of data sources, data collection procedures, and data collection. Then examined findings for transferability to ensure that the research findings were thorough, precise, organized, and correct. Then, we undertook the dependability test by auditing the entire study process. Thus, special might conduct confirmability tests to strengthen the objectivity of the research (Moran-Ellis et al., 2006). The observation step entails making direct observations of data sources in the four traditional villages of eastern Priangan in the form of flora present in the surrounding environment (Sugiyono, 2017). Interviews were done utilizing a semi-structured technique with informants comprising traditional leaders, indigenous peoples, and the younger generation who live in the four traditional villages of eastern Priangan (Sugiyono, 2017). The interview sample was chosen using a technique known as purposive sampling (Creswell and Creswell, 2018). They consider that the informants in the model are familiar with the use of edible plants as salads in East Priangan's four traditional villages. Additionally, researchers collected plant samples for laboratory morphological identification.

We carried out data analysis techniques, including data reduction, presentation, verification. Data reduction is made by summarizing, selecting the main parts, focusing on the essential things, looking for themes and patterns, and discarding unnecessary ones. After the data is reduced, the next step is to present the data in this study; the information is presented in tables, pictures, and narrative text. Finally, according to Miles and Huberman in Sugiyono (2015), the third step in qualitative data analysis is drawing conclusions and verification. And then, we will present the findings in this study after the data analysis stage has been completed and the data collected has been saturated. Then, to show the local importance of each species, Relative Frequency of Citation (RFC) analysis was carried out, and to determine a quantitative index to evaluate the relative usefulness of plants in an area, conducted use value (UV) analysis (Tardío and Pardo, 2008; Antayana, 2008; Ahmad et al., 2020; Bahadur et al., 2020). Relative frequency of citation (RFC) (Bahadur et al., 2020), which shows the local importance of each species, results from the equation:

$$RFC = \frac{Fc}{N}$$

F_c is the number of informants who mentioned the use of the species; N is the number of informants. The RFC has a value of 0–1.

Use value (UV) is a quantitative index to evaluate the relative usefulness in an area. It helps show which plants are most widely used to treat diseases in that area. High UV values indicate the main types of plants that are used as medicinal plants (Tardío and Pardo-de-Santayana, 2008; Fatima et al., 2018; Yaseen et al., 2019; Hussain et al., 2021). The following formula can calculate use value (UV):

$$UV = \frac{\sum U_i}{N}$$

U_i denotes the specific use of a plant species; N is the number of informants involved.

We also conducted correlation analysis to discover the correlation between the level of knowledge about indigenous vegetables and the age of indigenous people in indigenous villages in the East Priangan region (Chen and Popovich, 2011). To know the prospects of indigenous vegetables, an analysis using VOSviewer includes determining keywords, searching data in databases, selecting articles, validating data, and analyzing data (Eck and Waltman, 2010). Meanwhile, plant nomenclature was based on Flora Van Java (Backer and Bakhuizen Van Den Brink, 1968), with the author abbreviation following the Integrated Taxonomic Information System (ITIS, 2012).

3. Result and discussion

Indigenous knowledge found in indigenous village communities in East Priangan included local knowledge, understanding of customs and habits, nature, and how to build good relationships with all of them. Geographically, Kuta Village is located at the coordinates of 7° 16'13"S, 108° 33'48"E. Kuta Village is located separately from other villages because it is situated in a valley surrounded by perpendicular cliffs, which at the same time separate the village or become a boundary with other villages. The cliffs surrounding Kuta Village in the north, west, and south look like a fort that protects the village if you look from the inside of Kuta Village. As a valley area, Kuta Village is a fertile area with hilly and valley relief forms.

Meanwhile, Dukuh Village is at the coordinates of 7° 33'80"S, 107° 41'762"E. The topography of this traditional village is a valley

bordering the Cimangke River to the east and the Cisarangan River to the west; to the north, the village is bordered by Tutupan Forest and Titipan Forest. Pulo Village is located at the coordinates of 7° 6'11"S, 107° 55'7"E, on a small island that extends from west to east with 16.5 ha. Naga Village is in a fertile valley flanked by hills and is traversed by a river that has spring on the mountains, with coordinates 7° 21'42"S, 107° 59'35"E. In terms of topography, Naga Village has highlands and lowlands.

Nearly 72 percent of all indigenous people in East Priangan belong to a lower-middle-income group. They live depending on the agricultural sector as their primary source of income. The location of the traditional villages near or directly adjacent to the forest and rice fields is strongly tied to their employment as a farmer. The following is a general description of the area and community of Naga Village, which is presented in Fig. 1.

Edible plants that are consumed as lalapan are broadly divided into two groups. The first group is a type of vegetable deliberately planted as the property boundary, living fence, landslide barrier, green manure, ornamental plant, or medicinal plants in the garden. The second group is indigenous vegetables consumed as lalapan that grow wild, including plants that grow in the wild, such as mountains or forests, rice fields, gardens, or grasslands.

Sujarwo et al. (2016a) classified edible plants into wild edible plants, semi-wild edible plants, and cultivated edible plants. The status of wild plants changed to cultivated plants when the indigenous villagers deliberately planted them in their house yards. The good practice of local people in the four traditional villages in East Priangan is cooperation in a family manner. Each family makes it a habit of planting different types of plants in the garden or yard to complement the diverse needs of plants. The local people in the traditional villages also make it a habit to share information when they have experiences obtained by trial and error, such as experiences regarding processing methods, benefits, and specific recipes of herbal concoctions. This habit is passed down from generation to generation from their ancestors. The knowledge of plant usage, on the other hand, emerged in tandem with human civilization as a result of generations of hands-on experience with plants (Jan et al., 2011). Based on the interviews, the majority of informants had a positive response to indigenous vegetables in the form of wild plants that can be eaten




Fig. 1. Map of the area of Pulo Village, Kuta Village, Naga Village, and Dukuh Village in Priangan Timur, West Java.

as healthy and safe food, which is part of traditional food culture. Traditional knowledge in East Priangan is similar to Kaghlan Valley; Mansehra, Pakistan, believes that plants are essential for their life (Jamal et al., 2012).

3.1. Potency

The data collected from the four traditional villages, Pulo Village, Kuta Village, Naga Village, and Dukuh village, are presented in Table 1. Identified the collected specimen data morphologically to determine some of the characteristics of the specimens, including family, species name, local name, edible parts of plant organs, its benefits based on the local knowledge of indigenous people, how to eat it, whether it can be eaten fresh or requires prior processing, habitus, the status of the plant whether it is obtained from wild or has been cultivated by local people, and habitat type. The use-value (UV) and RFC (Relative Frequency of Citation) were calculated from these results.

The collection of data on edible indigenous vegetables consumed as lalapan at the four traditional villages showed the result of 86 species belonging to 32 families. Five species were intensively consumed as lalapan by the local people of the traditional villages in East Priangan based on the RFC value. The five plants were bonteng or cucumber (*Cucumis sativus* L.), whose fruit was consumed with RFC value = 0.90; sampeu or cassava (*Manihot esculenta* Crantz) whose leaves were consumed with RFC value = 0.86; papaya (*Carica papaya* L.) with RFC value = 0.83; genjer or yellow velvetleaf (*Limnorchis flava* L.) with RFC value = 0.71; and surawung or basil (*Ocimum basilicum* L.) with RFC value = 0.61. Some traditional vegetables had no clear definition due to their ambiguity. It was rather difficult to count and include them in the list of existing indigenous vegetables.

The edible indigenous plants' vegetable data are consumed as lalapan at the four traditional villages are similar to previous study in indigenous communities in Pakistan (Shaheen et al., 2017; Malik et al., 2018; Bahadur et al., 2020; Tariq et al., 2020). Meanwhile, in Pakistan, the plants use as medicinal treatment such as to treat musculoskeletal disorders (Malik et al., 2018a), hypertension treatment (Malik et al., 2018b), kidney stones treatment (Ahmad et al., 2018a), diabetes (Najab and Mushtaq, 2014; Abidin et al., 2018), herbal drinks (Rashid et al., 2018) or cosmetics (Ahmad et al., 2008) among indigenous communities in Pakistan. In Pakistan, edible plants and weeds are used for medicinal treatment, such as Convolvulaceae (Ashfaq et al., 2018). It is shown that indigenous plants benefit from indigenous knowledge in different communities.

Family classification of edible indigenous vegetables consumed as lalapan in four traditional villages in East Priangan can be seen in Fig. 2. Most of the edible indigenous vegetables consumed as lalapan were from the family Fabaceae, as many as ten species (13%). Nkonki et al. (2013) explained that the species of family Fabaceae are primarily nuts, which are the most nutritious food source. In other scientific studies, it turns out that this family produces a high diversity of secondary metabolites that function as compounds that avert herbivores and microbes and also as signaling compounds to attract pollinators and fruit-eating animals (Wink, 2013). Without them knowing the definition of secondary metabolites, the habit of eating fresh vegetables requires local people to cultivate edible plants as lalapan by planting them around the house or the village to meet their daily food needs.

Meanwhile, the least utilized families included Acanthaceae, Alismaceae, Amaranthaceae, Arecaceae, Portulacaceae, Caricaceae, Gnetaceae, Lecythidaceae, Moringaceae, Musaceae, Primulaceae, Pontederiaceae, Talinaceae, Theaceae, Woodsiaceae, and Xanthorrhoeaceae. The percentage of the use of vegetables from this families was 1%. Only a few people, generally adults and the elderly, ate lalapan from these families. The main reason was to meet daily needs as the primary food supplement and for other benefits such as herbal medicine or other health benefits.

Some parts of the plant that can be used as lalapan include young shoots and leaves, flowers and whorls, raw fruit, ripe fruit and seeds, tuber, bark, and skin of tuber. Lalapan in leaves (especially young leaves) ranked the most, with 62 species. The findings were in line with research by Cahyanto et al. (2018) on the Sundanese people in Subang Regency that at least people used 50 types of plants as lalapan, and 33 types of them were consumed raw or without going through the cooking process first. Furthermore, from 50 types of plants, the leaves of 32 of them were used for lalapan, which are consistently consumed with daily food.

The use of leaf parts of plants is indirectly based on the fact that the leaves of plants used as lalapan contain more essential elements such as vitamin B9, which plays a role in cell growth, prevents anemia, and helps the formation of red blood cells. Apart from containing vitamin B9, green vegetables also contain sufficient amounts of vitamin K, which plays a role in blood coagulation and prevents bleeding. Then, green vegetables contain other nutrients such as inositol (which plays a role in protecting the liver, heart, and digestive system), carotenoids, and provitamins that maintain the health of cells and body tissues (Butnariu and Butu, 2015).

Meanwhile, parts of edible plants used as lalapan include fruit, leaves, and nuts (Madison, 2008). In the Women Jungle Survival Course (2019), the parts of edible plants that can consume generally include leaves and young stems or stalks, which are usually greener and softer. Older plants are generally more bitter and rough. Roots and tubers are recommended to be selected from the larger plant. Especially in vines, there is a part in the root stem that can eat after tasting it first. It is advisable to choose fruits, nuts, seeds, and grains that are ripe, soft, contain less sap, and are bright in color.

3.2. Tradition

Classically, indigenous villagers in Priangan Timur, West Java, are familiar with lalapan (Amrinanto et al., 2019b; Hendariningrum, 2018). Their ancestors passed down the tradition of eating fresh vegetables since time immemorial. From generation to generation, the life of this classical community has a close interaction with resources and the environment, inheriting the traditional lifestyle that their ancestors lived. This conventional lifestyle then forms traditional wisdom.

Traditional wisdom is identified with human habits, nature, and the relationship between all inhabitants of the ecological community (Jaiswal, 2019; Makondo and Thomas, 2018). Consumption of plant species as lalapan is also identified with those three things. The indigenous villages know lalapan in the form of fresh vegetables eaten raw or cooked first to complement rice, side dishes, and sambal in their daily eating habits. Sambal is a sauce made from a mixture of chilies, tomatoes, galangal, garlic, shallots, sugar, salt, and shrimp paste. An example of a dish of lalapan with sambal is presented in Fig. 3.

Apart from being eaten raw with sambal, indigenous people in East Priangan are also accustomed to eating fresh vegetables as pecel (Hendariningrum, 2018). Pecel (Fig. 4a) is fresh vegetables (lalapan) that are boiled first, then mixed with peanut sauce that has been seasoned with spices like galangal. Sometimes, the term 'pecel' is replaced with 'karedok' (Fig. 4b), when the cooked lalapan are mixed in chopped raw lalapan. An example of a dish of lalapan as pecel is presented in Fig. 4. The concept of indigenous vegetables is also applied in several countries with dishes that are made to be appetizing, which differ in preparation techniques such as frying, making soup, or boiling for a long time (Luczaj et al., 2013).

Consumption of raw vegetables is synonymous with salads. Consuming salads shows a close relationship with the concept of diet, so it is very promising to increase the consumption of salad as part of the diet (Pieroni et al., 2015). Lalapan benefits health and skin because it contains nutrients, including β -carotene (Amrinanto et al., 2019a). Even Martins et al. (2011) explained that edible vegetables are nutritionally balanced.

1
42
Table 1
List of Potentially *lalapan* vegetables in Kampung Adat Priangan Timur.

Number	Family/ Species	Local Name	Parts used	Usability	Application	Habitus	Wild / Cultivated Status	Habitat Type	UV	1 RFC
1.	Acanthaceae <i>Staurogyne elongata</i> Luntze	Reundeu	FOL	hypertension medication	ED	HR	WL	MT; GD; FR; YRD; FR	1	0.33
2.	Alismataceae <i>Limncharis flava</i> L.	Genjer	FOL; CAU	cholesterol medication	B; ED	HR	WL	PI; FR	0.5	0.71
3.	Amaranthaceae <i>Amaranthus</i> <i>hybridus</i> L.	Bayem	FOL; CAU		B; S	HR	CV	GD; YRD; FR		0.27
4.	Anacardiaceae <i>Anacardium occiden- tale</i> L.	Jambu mede	FOL	Stomachache Medi- cine; Medicine for vaginal discharge; Shrinking wom- en's intimacy	ED	TR	CV	GD; YRD; FR	1.5	0.33
5.	<i>Mangifera indica</i> L.	Mangga	FOL		ED	TR	CV	YRD; SRD; GD		0.02
6.	<i>Bouea macrophylla</i> Griff.	Gandaria	FOL		ED	TR	CV	FR; GD; FR		0.02
7.	<i>Mangifera kemanga</i>	Kemang	FOL		ED	TR	CV	FR		0.02
8.	<i>Spondias dulcis</i> Parkinson	Kadongdong	FOL		B	TR	CV	YRD		0.13
9.	Apiaceae <i>Centella asiatica</i> (L.) Urban	Antanan gondrong/ Kurawet galeung	FOL; CAU; RAD	For herbs; Blood booster medica- tion; Ulcer medi- cine; wound medicine; Reduces choles- terol; Rheuma- tism medicine	ED	HR	WL	FR; MT; GD	0.5	0.49
10.	<i>Monochoria vaginalis</i> (Burm. f.) C. Presl ex Kunth	Antanan alit/ antanen beurit	FOL; CAU; RAD	treat jaundice; liver with edema; uni- nary tract infec- tions; coughing; shortness of breath; mouth sores, sore throat; and ear infections	ED	HR	WL	FR; MT; GD	0.5	0.49
11.	<i>Daucus carota</i> L.	Wortel	RAD	eye medicine	ED; B	HR	CV	FR; GD	0.5	0.07
12.	<i>Eryngium foetidum</i> L.	Walang geni/ Walang sangit	FOL	To lower choles- terol; Lowering high blood pressure	ED; B; S	HR	WL	YRD; FR; GD	2	0.07
13.	<i>Oenanthe javanica</i> DC.	Tespong	FOL		ED	HR	WL	PK, FR		0.12
14.	Arecaceae <i>Cocos nucifera</i> L.	Umbut Kalapa	CAU		B	TR	CV	FR		0.02
15.	Araliaceae <i>Hydrocotyle sibthor- pioides</i> Lamb.	Antanan alit	FOL	Blood booster	ED	HR	WL	FR	1	0.03
16.	<i>Polyscias fruticosa</i> (L.) Harms	Kadongdong cina/ Imba	FOL	For body aches; Increase appetite	ED; B	PR	CV	GD; SRD; YRD	0.6	0.11
17.	<i>Polyscias scutellaria</i> (Burm.f)	Mamangkakan	FOL		B; S	PR	CV	YRD; SRD; GD		0.05
18.	Asteraceae <i>Conyza sumatrensis</i>	Jalantir	FOL	Wound medicine	ED	PR	WL	FR; GD	1	0.13
19.	<i>Emilia sonchifolia</i> DC.	Jonge	FOL		ED	HR	WL	GD; FR		0.33
20.	<i>Lactuca sativa</i> L.	Saladah bokor	FOL		ED	HR	CV	YRD; GD; FR		0.11
21.	<i>Cosmos coudatus</i> Kunth.	Randa midang	FOL	Drugs for vaginal discharge	ED; B	PR	CV	YRD; SRD; GD	1	0.07
22.	<i>Gynura divaricata</i> L.	Seupan lentah	FOL		ED	HR	WL	FR		0.03
23.	<i>Bidens pilosa</i> L.	Ketul	FOL	Health potion	ED; B	HR	WL	GD; MT		0.02
24.	<i>Spilanthes paniculata</i> Wall. ex DC.	Jotang	FOL	To warm the throat	ED	HR	WL	YRD; GD; MT; FR; SRD	1	0.17
25.	<i>Crassocephalum</i> <i>crepidioides</i> (Benth.) S. Moore	Sintrong/Bagelis	FOL	Cholesterol drugs	ED	HR	WL	GD; YRD	1	0.21
26.	<i>Pluchea indica</i> (L.)	Baluntas	FOL	Drugs for vaginal discharge	ED; B	PR	WL	GD; YRD	0.5	0.06
27.	1 Brassicaceae <i>Brassica juncea</i> L.	Sawi	FOL		B	HR	CV	FR; GD		0.07

(continued)

Table 1 (Continued)

Number	Family/ Species	Local Name	Parts used	Usability	Application	Habitus	Wild / Cultivated Status	Habitat Type	UV	RFC
28.	<i>Brassica juncea</i> var 54 <i>gosa</i> L.	Jabung	FOL		ED; B	HR	CV	GD		0.03
29.	<i>Brassica oleracea</i> L.	Kol	8 FOL		ED; B	HR	CV	FR; GD		0.19
30.	<i>Brassica pekinensis</i> L.	Pecai	FOL		B	HR	CV	FR; GD		0.15
31.	<i>Brassica rapa</i> L.	Sausin	FOL		ED; B	HR	CV	FR		0.29
32.	<i>Raphanus sativus</i> L. Caricaceae	Lobak	RAD		ED; B	HR	CV	FR		0.02
33.	<i>Carica papaya</i> L.	Gedang	FOL; FR; FLO	Lowering high blood pressure; Increase appetite; Blood booster; Dengue medicine; Pain medicine for the body; Smooth digestion	B	PR	CV	GD; YRD; FR	0.75	0.83
34.	Convolvulaceae <i>Ipomoea batatas</i> (L.) Lam	Boled/Huwi	FOL	Raise platelets; Cho- lesterol drugs; Heat medicine; Hot / burn wound medicine	B; S	HR	CV	YRD; GD; FR	4	0.12
35.	<i>Ipomoea aquatic</i> Forsk.	Kangkung	FOL; CAU	Improves digestion; Sleep; Blood supplement	ED; B	HR	CV	PL; GD; FR	1.5	0.45
36.	Commelinaceae <i>Commelina bengha- lensis</i> L.	Gewor	FOL; CAU; RAD		B	HR	WL	GD; RF; SRD		0.02
37.	Cucurbitaceae <i>Cucumis sativus</i> L.	Bonteng	FRU	hypertension medication	ED	LN	CV	GD; FR; SRD; YRD	0.25	0.90
38.	40 <i>Cucurbita moschata</i> Duchesne	Waluh gede	FRU; FOL	hypertension medication	ED; B; S	39 LN	CV	FR; GD; YRD	0.5	0.45
39.	<i>Lagenaria siceraria</i> (Molina) Standl.	Leor	FRU	Fever medicine	B	LN	CV	GD; YRD	1	0.02
40.	<i>Luffa acutangula</i> (L.) Roxb	Emes (oyong)	FRU		B	LN	CV	GD; FR		0.02
41.	<i>Momordica charan- tica</i> Linn.	Paria/Pare	FRU; FOL	Medicine for headache	B	LN	CV	FR; MT; GD	1	0.43
42.	Euphorbiaceae <i>Cnidioscolus aconiti- folius</i> (Mill.) I.M. Johnst	Chaya-chaya	FOL		B; S	PR	CV	GD; YRD; SRD		0.06
43.	<i>Euphorbia hirta</i> L.	Nanangkaan	FOL	Drugs for vaginal discharge	ED	HR	WL	GD	1	0.02
44.	<i>Glochidion arbores- cens</i> Blume.	Mareme	FOL		ED	TR	CV	YRD; FR; GD		0.03
45.	<i>Manihot esculenta</i> Crantz	Sampeu	FOL	Blood booste; Ulcer medicine; Medi- cine for paralyzed pain; Wound medicine	B	PR	CV	GD; YRD; FR	0.2	0.86
46.	<i>Sauropus androgy- nus</i> (L.) Merr	Katuk	FOL	Eye medicine; Heat medicine; Streamlining breast milk	ED; B; S	PR	CV	YRD; GD; FR	0.3	0.29
47.	Fabaceae <i>Sesbania grandiflora</i> 48 Pers.	Kembang Turi	FLO		B; S	PR	WL	GD; SRD		0.03
48.	<i>Cajanus cajan</i> (L.) Huth	Hiris	FRU		ED	PR	CV	GD		0.03
49.	<i>Canavalia ensiformis</i> (L.) DC.	Koas	FRU		ED	HR	CV	GD		0.02
50.	<i>Leucaena leucoce- phala</i> (Lam.) de Wit	Peuteuy selong	FRU, FOL		ED	TR	CV	GD; YRD; SRD		0.24
51.	<i>Parkia javanica</i>	Peuteuy	FRU	Lowers cholesterol; Lowering high blood pressure	ED	TR	CV	GD; YRD	0.6	0.47
52.	<i>Pithecellobium loba- tum</i> Benth.	Jengkol	FRU	Dry sugar medicine; Lowers cholesterol	ED; B	TR	CV	GD; YRD	0.6	0.62
53.	<i>Phaseolus lunatus</i> L.	Roay/Kratok	FRU		ED; B	7 LN	CV	GD; YRD		0.17

(continued)

Table 1 (Continued)

Number	Family/ Species	Local Name	Parts used	Usability	Application	Habitus	Wild / Cultivated Status	Habitat Type	UV	RFC
54.	<i>Phaseolus vulgaris</i> L.	Buncis	FRU		ED; B	LN	CV	GD; FR		0.09
55.	<i>Psophocarpus tetragonolobus</i> (L.) DC.	Jaat/Kecipir	FRU		ED; B	LN	CV	GD; FR; SRD		0.13
56.	<i>Vigna cylindrica</i> (L.) Skeels. Gnetaceae	Kacang panjang/ Kacang turus	FRU; FOL		ED; B; S	LN	CV	SRD; FR; GD; YRD		0.58
57.	<i>Gnetum genom</i> L. Lamiaceae	Tangkil	FOL		B	TR	CV	GD		0.02
58.	<i>Ocimum basilicum</i> L.	Surawung	FOL		ED	PR	CV	YRD; GD; FR		0.61
59.	<i>Plectranthus amboinicus</i> (Lour.) Spreng Lecythidaceae	Jinten	FOL		ED; B	HR	CV	GD; YRD		0.09
60.	<i>Planchonia valida</i> Bl.	Putat/Jaha	FOL	Bloating medicine; Getting close to a woman's inti- macy; Diarrhea medicine	ED	TR	WL	MT; GD; FR; YRD; FR	0.6	0.45
61.	Moraceae <i>Artocarpus hetero- phyllus</i> Lam	Nangka (tongtolang)	FRU		B	TR	CV	GD		0.09
62.	<i>Ficus fistulosa</i> Reinw ex. Blume	Beunying	FOL		ED	TR	WL	GD; FR		0.02
63.	<i>Ficus sundaica</i> Blume. Moringaceae	Koang	FOL	Diarrhea medicine	B	TR	WL	FR; GD	1	0.03
64.	<i>Moringa oleifera</i> L.	Kelor	FOL	Cure all diseases; Cholesterol drugs; Gout medicine; Rheumatism medicine	B; S	TR	CV	GD	2	0.03
65.	Musaceae <i>Musa paradisiaca</i> L.	Ower /Jantung pisang	SPAT		ED; B	HR	CV	GD		0.02
66.	56 eae <i>Bambusa vulgaris</i> Schrader ex J.C. Wendl	Iwung/Rebung	CAU		B	PR	WL	FR; GD		0.05
67.	<i>Zea mays</i> L.	Putri jagong	FRU		ED; B	HR	CV	FR; GD		0.03
68.	51 ederiaceae <i>Monochoria vaginalis</i> (Burm. f.) C. Presl ex Kunth	Eceng leutik/ Wewehan	FLO; CAU; FOL		B	HR	WL	RF; YRD; PL		0.02
69.	Portulacaceae <i>Portulaca oleracea</i> L.	Krokot	FOL; CAU; RAD		B; S	HR	WL	GD; RF		0.02
70.	Primulaceae <i>Ardisia elliptica</i> Thunb.	Ki lampeuni	FOL		ED	TR	WL	GD; FR		0.03
71.	Rubiaceae <i>Paederia foetida</i> L.	Kahitutan	8 FOL	Bloating medication	ED; B	LN	47 CV	SRD; GD; YRD	0.25	0.12
72.	<i>Morinda citrifolia</i> L.	Mengkudu	FOL	Pain medicine for the body	B	TR	CV	YRD; GD	1	0.03
73.	Solanaceae <i>Solanum lycopersi- cum</i> L.	Kemir	FRU		ED	PR	CV	YRD; FR; GD		0.05
74.	<i>Solanum macrocar- pon</i> L.	Terong buweuk	FRU		ED; B	PR	CV	GD; YRD		0.03
75.	<i>Solanum melongena</i> L.	Terong gelatik	FRU		ED; B	PR	CV	GD; YRD		0.53
76.	<i>Solanum nigrum</i> L.	Leunca	FRU; FOL		ED	PR	CV	YRD; GD; FR		0.49
77.	<i>Solanum torvum</i> Swartz Talinaceae	Takokak/magae	FRU	Diarrhea medicine; Itching medicine	ED; B	PR	WL	GD; SRD	1	0.17
78.	<i>Talinum paniculatum</i> (Jacq.) Gaertn	Ginseng jawa	FOL; CAU	Pain medicine for the body	ED; B	HR	CV	GD; YRD	1	0.02
79.	Urticaceae <i>Pilea trinervia</i> Wight.	Pohpohan	FOL		ED	HR	WL	MT; FR; GD; FR		0.25
80.	<i>Villebrunea rubes- cens</i> (Bl.) Bl.	Nangsi	FOL		ED	TR	WL	GD; FR		0.02
81.	Woodsiaceae <i>Diplazium esculen- tum</i> (Retz.) Sw.	Pakis sayur	FOL		B; S	HR	WL	MT; FR; FR		0.03
82.	Xanthorrhoeaceae <i>Aloe vera</i> Burm. f.	Lidah buaya	FOL		B	HR	CV	GD; YRD	2	0.02

(continued)

Table 1 (Continued)

Number	Family / Species	Local Name	Parts used	Usability	Application	Habitus	Wild / Cultivated Status	Habitat Type	UV	RFC
				treat heartburn; hair fertility						
83.	Zingiberaceae <i>Alpinia galangan</i> L.	Laja	RHI	Pain medicine	ED	HR	CV	GD; YRD	1	0.02
84.	<i>Curcuma longa</i> L.	Koneng	RHI	Ulcer medication / lowering stomach acid	ED	HR	CV	YRD; GD	0.25	0.15
85.	<i>Etlingera elatior</i> (Jack)	Comrang/Rombeh	CAU		ED	HR	CV	GD		0.03
86.	<i>Kaempferia galanga</i> L.	Cikur	FOL	Swelling medicine	ED; B	HR	CV	YRD; GD	1	0.09

Description

Parts used: CAU: caulis (stem); RHI: rhizoma (rhizome); RAD: radix (root); FOL: folium (leaves); FLO: flos (flower); FRU: fructus (fruit);.

SPAT: spatha (sheath).

Application: ED: eaten directly; B: boiled; S: steamed.

Habitus: TR: tree; PR: perdu; LN: liana; H: herbs.

Habitat Type: MT: mountain; FR: forest; RF: rice fields; YRD: yard; GD: garden; PL: Pool; SRD: side of the road.

Wild/Cultivate : CV: cultivation; WL: wild.




Fig. 2. Familia of edible indigenous vegetables that are consumed as lalapan in four traditional villages in East Priangan.

3.3. Local knowledge

The level of education of the local people at traditional villages in Priangan Timur was dominated by the elementary school level, so the people are considered to have a low level of education. However, the level of education of these indigenous people did not significantly

influence the level of knowledge about edible and medicinal plants and their farming activities. The local genius of the indigenous people is obtained from their ancestors based on experiences. The tradition of eating indigenous vegetables has declined over the years.

Lalapan tended to be better known by women with an average age above 50 years. The well-known and perceived benefit stimulus



Fig. 3. Lalapan with Sambal as Sauce.



Fig. 4. Lalapan is served as pecel (A) and karedok (B).

is that some fresh vegetables have benefits as traditional medicines, which local people believe treat more than one disease. The correlation between the age of the informants in the traditional village and the level of knowledge about lalapan at four traditional villages in East Priangan has analyzed the value of α 0.05 (5%), as presented in Table 2.

Table 2 shows a significance value of 0.000, which was smaller than the value of α 0.05 and meant that there was a positive correlation between age and the people's level of knowledge about indigenous vegetables that were consumed as lalapan. The correlation coefficient (R) was 0.587, which was in the excellent category, and the coefficient of determination (R-squared) was 0.344 or equal to 34.4%. This coefficient meant that the age of the people contributed to the knowledge level variable by 34.4%. In comparison, the

remaining 65.4% was the contribution of other variables that did not examine in this study. The contribution of these different variables was an error in this study. However, the results in Table 2 found that the coefficient of determination (R-Squared) was lower than the value of the contribution of other variables. Therefore, we can say that the contribution was low. The linear regression line is shown by a scatterplot (bivar) = X with Y in Fig. 5.

Based on the scatterplot presented in Fig. 5, a linearity line pattern can be seen between age and people's level of knowledge about indigenous vegetables consumed as lalapan. The dots on the plot indicate the amount of error from the correlation. The regression equation obtained from the analysis was $\hat{Y} = 9 + 0.1 * x$. The value of a was positive, meaning that if the age variable did not exist or were equal to zero, the level of knowledge would decrease. If the age were

33 e2
The results of the analysis of the correlation between age and the level of knowledge.

Mode	R	R-Squared	Adjusted R-Squared	Std. Error of the Estimate	Change Statistics					Sig. F Change
					R Square Change	F Change	df1	df2		
1	.587 ^a	.344	.330	1.182	.344	24.647	1	47		.000




Fig. 5. Scatterplot with Regression Line.

increased by 1, the level of knowledge would increase by 0.1. This positive correlation could mean that if the age of the indigenous people of traditional villages were getting older, the level of knowledge would also increase. The results of the research conducted by Uchiyama et al. (2017) showed that the level of recognition of traditional vegetables correlated with the age of the respondents.

3.4. Future prospect

The use of plants in people's lives can be seen in tri-stimuli, namely, natural stimulus, beneficial stimulus, and religious stimulus (Zuhud et al., 2007). Natural stimulus in the form of people's innate knowledge about plants, valuable stimulus related to the benefits or interests of the people towards plants, and religious or spiritual stimulus is the willingness and morals of the people to take conservation actions.

The correlation between utilization and people related to the role of the stimulus leads to the interest in sustainable use. However, sometimes, indigenous knowledge is undermined quickly. This loss of expertise negatively affects the lives and health of rural communities who live traditionally. For this reason, the preservation of indigenous knowledge is essential. Dweba & Mearns (2011) argued that the transfer of indigenous knowledge about traditional vegetables would ensure the availability and utilization of this critical food source for resource-poor communities. Chen & Qiu (2012) explained that the extraction of non-timber forest products is a way to create healthy socioeconomic relationships that depend on the surrounding biodiversity. The interaction of people with existing natural resources requires them to maintain indigenous knowledge to live in harmony with nature.

Indigenous vegetables are used as lalapan by the people in the four traditional villages and are not sold to people outside the traditional villages. They only plant on a small scale for personal consumption or share with neighbors in need. Most fresh vegetables and edible plant species with secondary functions as traditional medicines are grown and cultivated in gardens and yards.

Planting beneficial species in the yard is based on their function, for example, as a source of carbohydrates, protein, vitamins and minerals, medicinal plants, and others. This action is a form of conservation conducted by the local people that leads to natural stimuli, namely, the cultivation of edible plants consumed as lalapan. Local

people only cultivate plants that are needed in their daily life. Meanwhile, the religious stimulus in terms of sustainable use of natural resources is a self-sacrificing attitude to realize conserving natural resources in the vicinity. A participatory approach by involving local communities in plant conservation activities is an excellent supporting factor (Ulian et al., 2017).

Scholars and stakeholders are increasingly attaching the importance to local wild foods to evaluate and assess local biocultural heritage (Pieroni et al., 2015). For prospects, it is crucial to make an effort to obtain the primary data that are needed for the preservation of potential plant use, both in terms of bioecology and cultivation techniques and exploration of beneficial active compounds. Another function related to the sustainability of indigenous vegetables is to protect flora (Uchiyama et al., 2017).

To discover the extent to which research on indigenous vegetables has been carried out to prospects around the world and to know research trends and research collaborations, We made a bibliometric map of science. Bibliometric indicators are knowledge that can be used to evaluate scientific research results. It is widely used to produce a mapping of fields of science and to track or trace the development of new knowledge in a particular area and is used as an indicator in the future in providing a more competitive advantage and in making strategic plans (Mallig, 2010; Leydesdorff and Rafols, 2012).

To obtain data on indigenous vegetables about prospects, We searched an international database, Scopus, for 2015–2020. The search results selected the collected journal articles to ensure that the articles matched the topics and keywords used. The search results obtained for articles that fit the topic were 398 articles. The data were obtained using the keywords of 'indigenous vegetable' and 'prospects'. The network visualization display is presented in Fig. 6.

Fig. 6 shows that reduced the research trend on indigenous vegetables about prospects to 4 clusters. The advantage of VOSviewer over other analytical applications is that it uses a text-mining function to identify combinations of noun phrases relevant to mapping and an integrated clustering approach to examine data co-citation networks and co-occurrence. Interactive options and functions make this program accessible and easy to explore bibliometric data networks, such as the number of citations and co-occurrence relationships between key terms and concepts (Hofmann and Chisholm, 2016; Miner et al., 2012).




Fig. 6. Visualization of network trend of research publication based on co-words.

45
The results of the bibliometric knowledge map showed that the prospects of research on indigenous vegetables are still broad. Indigenous people are indifferent clusters in many countries, such as China and India. Indigenous people in Indonesia still have many opportunities to explore more profoundly, especially in terms of sustainability.

4. Conclusion

Lalapan is an essential part of a food source. The experience of indigenous people in the East Priangan region can explain that the details of plants that can eat as lalapan included young shoots and leaves, flowers and whorls, raw fruit, ripe fruit and its seeds, tuber, bark, and skin of tuber. 44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We want to thank Universitas Siliwangi for funding this research through the Simpemaus Internal Research Grant as per decision number 266/ UN58.21/PG/2020 regarding the assignment of an international journal-oriented research program in 2020. We also say thank to Universitas Airlangga for helping to proof read our manuscript by Airlangga Editing Program.

Reference

- Abbasi, A.M., Khan, M.A., Shah, M.H., Shah, M.M., Pervez, A., Ahmad, M., 2013. Ethnobotanical appraisal and cultural values of medicinally important wild edible vegetables of lesser himalayas-Pakistan. *J. Ethnobiol. Ethnomed.* <https://doi.org/10.1186/1746-4269-9-66>.
- Abidin, S.Z.U., Khan, R., Ahmad, M., Zafar, M., Saeed, A., Khan, N., 2018. Ethnobotanical survey of highly effective medicinal plants and phytotherapies to treat diabetes mellitus II in South-West Pakistan. *Indian J. Traditional Med.* 17 (4), 682–690.
- Ahmad, M., Butt, M.A., Zhang, G., Sultana, S., Tariq, A., Zafar, M., 2018. *Bergenia ciliata*: a comprehensive review of its traditional uses, phytochemistry, pharmacology and safety. *Biomed. Pharmacother.* 97, 708–721.
- Ahmad, M., Khan, M.A., Zafar, M., 2008. Traditional herbal cosmetics used by local women communities in district Attock of Northern Pakistan. *Indian J. Traditional Knowledge* 7 (3), 421–424.
- Ahmad, M., Malik, K., Tariq, A., Zhang, G., Yaseen, G., Rashid, N., Sultana, S., Zafar, M., Ullah, K., Khan, M.P.Z., 2018. Botany, ethnomedicines, phytochemistry and pharmacology of Himalayan Paeony (*Paeonia emodi* Royle). *J. Ethnopharmacol.* 220, 197–219.
- Ahmad, S., Zafar, M., Shinwari, S., Ahmad, M., Shinwari, Z.K., Sultana, S., 2020. Ethnobotanical Plants And Traditional Knowledge Linked To Primary Health Care Among The Indigenous Communities Living In Western Hilly Slopes Of Dera Ghazi Khan, Pakistan. *Pak J. Bot.* 52 (2), 519–530.
- Alves, R., 2016. Relationships Between Fauna and People and the Role of Ethnobiology in Animal Conservation. *Ethnobiology and Conservation* August 2012. <https://doi.org/10.15451/ec2012-8-1.2-1-69>.
- Aminanto, A.H., Hardinsyah, H., Palupi, E., 2019. The eating culture of the Sundanese: does the traditional salad (Lalapan) improve vegetable intake and blood β -carotene concentration? *Future of Food: Journal on Food, Agriculture and Society*, 7(2 SE-Research Articles) Article Nr. 104. <https://doi.org/10.17170/kobra-20190709593>.
- Backer, C.A., Bakhuizen Van Den Brink, R.C., 1968. *Flora of Java*, 3. *Flora of Java*.
- Bahadur, S., Khan, M.S., Shah, M., Shuaib, M., Ahmad, M., Zafar, M., Begum, N., Gul, S., Ashfaq, S., Mujahid, I., Hussain, F., 2020. Traditional usage of medicinal plants among the local communities of Peshawar valley, Pakistan. *Acta Ecologica Sinica* 40 (1), 1–29.
- Bhatia, H., Sharma, Y.P., Manhas, R.K., Kumar, K., 2018. Traditionally used wild edible plants of district Udhampur, J&K, India. *J. Ethnobiol. Ethnomed.* <https://doi.org/10.1186/s13002-018-0272-1>.
- Buanasari, B., Eden, W.T., Sholichah, A.I., 2017. Extraction of Phenolic Compounds From Petal Leaves (*Parkia Speciosa* Hassk.) Using Microwave and Ultrasound Assisted Methods. *Jurnal Bahan Alam Terbarukan* 6 (1), 25–31. <https://doi.org/10.15294/jbat.v6i1.7793>.
- Butnariu, M., Butu, A., 2015. Chemical Composition of Vegetables and Their Products. *Handbook of Food Chemistry*. Springer-Verlag Berlin Heidelberg, pp. 1–49. <https://doi.org/10.1007/978-3-642-36605-5>. Issue Januari 2015.
- Cahyanto, T., Supriyatna, A., Sholikha, M., Saepuloh, A., Rahmawati, D., 2018. Inventory of Plants Used as Lalapan in Subang. The 9th International Conference on Global Resources Conservation (ICGRC) and AII from Ritsumeikan University, West Jav-app. 1–11. <https://doi.org/10.1063/1.5061843>.

- Cartea, M.E., Francisco, M., Soengas, P., Velasco, P., 2011. Phenolic compounds in Brassica vegetables. *Molecules* 16 (1), 251–280. <https://doi.org/10.3390/molecules16010251>.
- Chen, B., Qiu, Z., 2012. Consumers' Attitudes towards Edible Wild Plants: a Case Study of Noto Peninsula, Ishikawa Prefecture, Japan. *International Journal of Forestry Research*. <https://doi.org/10.1155/2012/872413>.
- Chen, P.Y., Popovich, P.M., 2011. Correlation: Parametric And Nonparametric Measures. *Correlation*. <https://doi.org/10.4135/9781412983808.n1>.
- Chhikara, N., Devi, H.R., Jaglan, S., Sharma, P., Gupta, P., Panghal, A., 2018. Bioactive compounds, food applications and health benefits of *Parkia speciosa* (stinky beans): a review. *Agriculture and Food Security* 7 (1), 1–9. <https://doi.org/10.1186/s40066-018-0197-x>.
- Cita, K.D., 2020. Ethnobotany of food plant used by Sundanese Ethnic in Kalapare Village, Nyangkewok Hamlet, Sukabumi District, Indonesia. *Asian Journal of Ethnobiology* 3 (1), 16–22. <https://doi.org/10.13057/asianjethnobiology030103>.
- Creswell, J.W., Creswell, J.D., 2018. *Research and Design Qualitative, Quantitative and Mixed Methods Approaches*. Thousand Oaks California.
- Dalar, A., Uzun, Y., Turker, M., Mukemre, M., Konczak, I., 2016. Health attributes of ethnic vegetables consumed in the Eastern Anatolia region of Turkey: antioxidant and enzyme-inhibitory properties. *Journal of Ethnic Foods*. <https://doi.org/10.1016/j.jef.2016.05.005>.
- Dewi, Z.L., Halim, M.S., & Derksen, J. (2017). *Emotional intelligence competences of three different ethnic groups in Indonesia*. 1369 (April). <https://doi.org/10.1080/14631369.2017.1310615>.
- Dweba, T.P., Mearns, M.A., 2011. Conserving indigenous knowledge as the key to the current and future use of traditional vegetables. *Int J Inf Manage*. <https://doi.org/10.1016/j.ijinfomgt.2011.02.009>.
- Fatima, A., Ahmad, M., Zafar, M., Yaseen, G., Khan, M.P.Z., Butt, M.A., Sultana, S., 2018. Ethnopharmacological relevance of medicinal plants used for the treatment of oral diseases in Central Punjab-Pakistan. *J Herb Med* 12, 88–110.
- Guarnera, P.M., Savo, V., 2016. Wild food plants used in traditional vegetable mixtures in Italy. *J Ethnopharmacol*. <https://doi.org/10.1016/j.jep.2016.02.050>.
- Guerrero, D.J.L.G., 2015. *Edible Wild Plants. Recent Progress in Medicinal Plants. January 2002*.
- Hadi, U.K., Takaoka, H., 2018. The biodiversity of black flies (Diptera: simuliidae) in Indonesia. *Acta Trop.* 185, 133–137. <https://doi.org/10.1016/j.actatropica.2018.02.013>.
- Harashani, H., 2018. Local Wisdom of Kampung Naga in The Era of Globalization. *Journal of Humanities and Social Studies* 2 (1), 51–54. <https://doi.org/10.33751/jhss.v2i1.823>.
- Hendariningrum, R., 2018. *Budaya dan Komunikasi Kesehatan (Studi Pandangan Kesehatan Pada Masyarakat Sunda Dalam Tradisi Makan Lalapan)*. LUGAS: Jurnal Komunikasi 2 (1), 13–19. <https://doi.org/10.31334/jlv2i1.118>.
- Hidayat, S., Hikmat, A., Zuhud, E.A.M., 2010. Kajian Etnobotani Masyarakat Kampung Adat Dukuh Kabupatan Garut, Jawa Barat. *Media Konservasi* 15 (3), 139–151.
- Hofmann, M., Chisholm, A., 2016. Text Mining and Visualization: Case Studies Using Open-Source Tools. *Text Mining and Visualization: Case Studies Using Open-Source Tools*.
- Hussain, A., Zafar, M., Shinwari, S., Shinwari, Z.K., Ahmad, M., Sultana, S., Yaseen, G., 2021. Ethnoverinary uses of medicinal plants as herbal drugs for sustainable livestock in southern deserts of Sindh Pakistan. *Pakistan Journal of Botany*, 53, 2.
- Iliyani, M., 2018. Thinking Intellectually, Knowing Globally, and Act Locally: the Prospect of Kampung Naga as An Indigenous Village. *Jurnal Masyarakat & Budaya* 20 (1), 15–30. <https://doi.org/10.14203/jmb.v20i1.588>.
- Integrated Taxonomic Information System. (2012). *Choice Reviews Online*. <https://doi.org/10.5860/choice.49-6871>.
- Jaiswal, A. (2019). *Indigenous Knowledge System and Traditional Medicinal Practitioner*. April. <https://doi.org/10.23880/aeaj-16000119>.
- Jamal, Z., Ahmad, M., Zafar, M., Sultana, S., Khan, M.A., Shah, G.M., 2012. Medicinal plants used in traditional folk recipes by the local communities of Kaghlan valley, Mansehra, Pakistan. *Indian J. Traditional Knowl.* 11 (4), 634–639.
- Jan, G., Khan, M.A., Ullah, F., Jan, F.G., Ahmad, M., Jan, M., Zafar, M., 2011. Ethnobotanical studies on some useful plants of Dir Kohistan valleys. *KPK, Pakistan. Pakistan Journal of Botany* 43 (4), 1849–1852.
- Kang, Y., Łuczaj, Ł., Kang, J., Wang, F., Hou, J., Guo, Q., 2014. Wild food plants used by the Tibetans of Gongba Valley (Zhouqu county, Gansu, China). *J. Ethnobiol. Ethnomed*. <https://doi.org/10.1186/1746-4269-10-20>.
- King, S.R., Carlson, T.J., Moran, K., 1996. Biological diversity, indigenous knowledge, drug discovery and intellectual property rights: creating reciprocity and maintaining relationships. *J. Ethnopharmacol.* 51 (1–3), 45–57. [https://doi.org/10.1016/0378-8741\(95\)01349-0](https://doi.org/10.1016/0378-8741(95)01349-0).
- Leydesdorff, L., Rafols, I., 2012. Interactive overlays: a new method for generating global journal maps from Web-of-Science data. *J. Informetr.* <https://doi.org/10.1016/j.joi.2011.11.003>.
- Lincoln, Y.S., Guba, E.G., 1985. *Naturalistic Inquiry*, 75th ed. Sage.
- Łuczaj, Ł., Zovko-Končić, M., Milčević, T., Dolina, K., Pandza, M., 2013. Wild vegetable mixes sold in the markets of Dalmatia (southern Croatia). *J. Ethnobiol. Ethnomed*. <https://doi.org/10.1186/1746-4269-9-2>.
- Madison, D. (2008). *Edible An Illustrated Guide to the World's Food Plants*.
- Majeed, S., Zafar, M., Ahmad, M., Ozdemir, F.A., Kilic, O., Hamza, M., Sultana, S., Yaseen, G., Raza, J., 2021. Ethnobotany, Medicinal Utilization and Systematics of *Opuntia* Species from Deserts of Pakistan. *Opuntia spp Chemistry, Bioactivity and Industrial Applications*. Springer, Cham, pp. 49–80.
- Makondo, C.C., Thomas, D.S.G., 2018. Climate Change adaptation: Linking indigenous Knowledge With Western Science For Effective Adaptation. *Environmental Science and Policy*. <https://doi.org/10.1016/j.envsci.2018.06.014>.
- Malik, K., Ahmad, M., Zhang, G., Rashid, N., Zafar, M., Sultana, S., Shah, S.N., 2018. Traditional plant-based medicines used to treat musculoskeletal disorders in Northern Pakistan. *Eur. J. Integr. Med* 19, 17–64.
- Alghatani Malik, K., Ahmad, M., Bussmann, R.W., Tariq, A., Ullah, R., Ali, S., Shahat, A.A., Rashid, N., Zafar, M., Sultana, S., Shah, S.N., 2018. Ethnobotany of Anti-hypertensive Plants Used in Northern Pakistan. *Front. Pharmacol* 09, 1–18.
- Mallig, N., 2010. A relational database for bibliometric analysis. *J. Informetr.* <https://doi.org/10.1016/j.joi.2010.06.007>.
- Martins, D., Barros, L., Carvalho, A.M., Ferreira, I.C.F.R., 2011. Nutritional and in Vitro Antioxidant Properties of Edible Wild Greens in Iberian Peninsula Traditional Diet. *Food Chemistry*. <https://doi.org/10.1016/j.foodchem.2010.09.038>.
- Miner, G.D., Elder, J., Nisbet, R.A., 2012. *Practical Text Mining and Statistical Analysis For Non-structured Text Data Applications. Practical Text Mining and Statistical Analysis for Non-structured Text Data Applications*. <https://doi.org/10.1016/C2010-0-66188-8>.
- Mirajab, K., Mushtaq, A., 2014. Potential for Pakistani traditional medicinal plants to combat diabetes. *J. Tradit. Chin. Med.* 34 (4), 488–490.
- Mokganya, M.G., Mushaphi, L.F., Tshisikhawe, M.P., 2019. Indigenous preparation and preservation methods of selected indigenous wild edible vegetables of the Vhembe district municipality. *Indilinga African J. Indigenous Knowledge Syst.*.
- Moran-Ellis, J., Alexander, V.D., Cronin, A., Dickinson, M., Fielding, J., Sleney, J., Thomas, H., 2006. Triangulation and integration: processes, claims and implications. *Qualitative Research* 6 (1), 45–59. <https://doi.org/10.1177/1468794106058870>.
- Ahmad, & M. Nazish, M., Zafar, M., Sultana, S., 2020. Halophyte Diversity in Pakistan" Wild Resources and their Ethnobotanical Significance. *Handbook of Halophytes*. Springer Nature Switzerland.
- Negi, V.S., Pathak, R., Sekar, K.C., Rawal, R.S., Bhatt, I.D., Dhyani, P.P., Negi, V.S., Pathak, R., Sekar, K.C., Rawal, R.S., & Bhatt, I.D. (2017). *Traditional knowledge and biodiversity conservation : a case study from Byans Valley in Kailash Sacred Landscape, India*. 0568(September). <https://doi.org/10.1080/09640568.2017.1371006>.
- Nkonki, T., Ngwenya, A.M., & Van Wyk, B.-E. (2013). *Fabaceae, Bothalia*. <https://doi.org/10.4102/abc.v4i2.102>.
- Padhy, S., Dash, S.K., Mohapatra, R., Padhy, S., Dash, S.K., Mohapatra, R., 2017. *Ethnobiology : the Science of a Common Man to Explore the Traditional Ecology, Biodiversity and Philosophy of Conservation : a Review Ethno-biology : the Science of a Common Man to Explore the Traditional Ecology*. Biodiversity and Philosophy of Conserv 6901. <https://doi.org/10.1080/09766901.2015.11884751>.
- Perchuk, I., Shelenga, T., Gurkina, M., Miroshnichenko, E., Burlyayeva, M., 2020. composition of primary and secondary metabolite compounds in seeds and pods of asparagus bean (*Vigna unguiculata* (L.) Walp.) from China. *Molecules* 25 (17), 1–16. <https://doi.org/10.3390/molecules25173778>.
- Pieroni, A., Nedelcheva, A., Dogan, Y., 2015. Local knowledge of medicinal plants and wild food plants among Tatars and Romanians in Dobruja (South-East Romania). *Genet. Resour. Crop Evol.* <https://doi.org/10.1007/s10722-014-0185-3>.
- Purba, S.F., Chahaya, I., Marsaulina, I., 2013. Pemeriksaan Escherichia coli dan Larva Cacing pada Sayuran Lalapan Kemangi (*Ocimum basilicum*), Kol (*Brassica oleracea* L. var. capitata L.), Selada (*Lactuca sativa* L.), Terong (*Solanum melongena*) yang Dijual di Pasar Tradisional, Supermarket, dan Restoran. *Jurnal Lingkungan Dan Kesejahteraan Kerja* 2 (1), 80–87. <https://doi.org/10.1016/B978-012373960-5.00477-9>.
- Qodariah, L., Armiyati, L., 2013. Nilai-Nilai Kearifan Lokal Masyarakat Adat Kampung Naga Sebagai Alternatif Sumber Belajar. *SOCIA: Jurnal Ilmu Sosial* 10 (1), 10–20. <https://doi.org/10.21831/socia.v10i1.5338>.
- Rashid, N., Gbedomon, R.C., Ahmad, M., Salako, V.K., Zafar, M., Malik, K., 2018. Traditional knowledge on herbal drinks among indigenous communities in Azad Jammu and Kashmir. *Pakistan J. Ethnobiol. Ethnomed.* 14 (1), 16–36.
- Saba, A., Moneta, E., Pepparo, M., Sinesio, F., Vassallo, M., Paoletti, F., 2018. Towards a Multi-Dimensional Concept of Vegetable Freshness from the Consumer's Perspective. *Food Quality and Preference*. <https://doi.org/10.1016/j.foodqual.2017.12.008>.
- Saleh, M.S.M., Jalil, J., Zainalabidin, S., Asmadi, A.Y., Mustafah, N.H., Kamisah, Y., 2021. Genus parkia: phytochemical, medicinal uses, and pharmacological properties. *Int J Mol Sci* 22 (2), 1–42. <https://doi.org/10.3390/ijms22020618>.
- Shaheen, S., Abbas, S., Hussain, J., Mabood, F., Umair, M., Ali, M., Ahmad, M., Zafar, M., Khan, U.F.A., 2017. Knowledge of medicinal plants for Children diseases in the environs of District Bannu, Khyber Pakhtoonkhwa (KPK). *Front. Pharmacol* 8, 430.
- Singhania, N., Chhikara, N., Bishnoi, S., Garg, M.K., Panghal, A., 2021. Bioactive Compounds of Petai Beans (*Parkia Speciosa* Hassk.). *Reference Series in Phytochemistry*, pp. 525–543. https://doi.org/10.1007/978-3-030-57415-4_30. October.
- Sudjatnika, T., 2018. Filosofi Hidup Komunitas Masyarakat Adat Sunda Kampung Naga Ditinjau dari Pranata Keagamaan. *Jurnal Al-Tsaqafa* 15 (1), 69–76. <https://doi.org/10.1017/CBO9781107415324.004>.
- Sugiyono, 2017. *Metode Penelitian Kuantitatif, Kualitatif, dan R&D*. Alfabeta.
- Sujarwo, W., Arinasa, I.B.K., Caneva, G., Guarnera, P.M., 2016a. Traditional knowledge of wild and semi-wild edible plants used in Bali (Indonesia) to maintain biological and cultural diversity. *Plant Biosyst.* <https://doi.org/10.1080/11263504.2014.994577>.
- Sujarwo, W., Caneva, G., 2016b. Using quantitative indices to evaluate the cultural importance of food and nutraceutical plants: comparative data from the Island of Bali (Indonesia). *J. Cult Herit* 18. <https://doi.org/10.1016/j.culher.2015.06.006>.
- Tardío, J., Pardo-de-Santayana, M., 2008. Cultural importance indices: a comparative analysis based on the useful wild plants of Southern Cantabria (Northern Spain). *Econ. Bot.* 62, 24–39.
- Tariq, A., Sadia, S., Fan, Y., Ali, S., Amber, R., Mussarat, S., Ahmad, M., Murad, W., Zafar, M., Adnan, M., 2020. Herbal medicines used to treat diabetes in Southern regions of Pakistan and their pharmacological evidence. *J Herb Med* 21, 100323.
- Thomas, E., Magilvy, J.K., 2011. Qualitative Rigor or Research Validity in Qualitative Research. *J. Spec. Pediatr. Nurs.* 16 (2), 151–155. <https://doi.org/10.1111/j.1744-6155.2011.00283.x>.

- Tufts, H.R., Harris, C.S., Bukania, Z.N., Johns, T., 2015. Antioxidant and Anti-Inflammatory Activities of Kenyan Leafy Green Vegetables, Wild Fruits, and Medicinal Plants with Potential Relevance for Kwashiorkor. *Evid.-Based Complement. Altern. Med.* <https://doi.org/10.1155/2015/807158>.
- Uchiyama, Y., Fujihira, Y., Matsuo, H., Kohsaka, R., 2017. Tradition and Japanese vegetables: history, locality, geography, and discursive ambiguity. *J. Ethnic Foods.* <https://doi.org/10.1016/j.jef.2017.08.004>.
- Ulian, T., Sacandé, M., Hudson, A., Mattana, E., 2017. Conservation of indigenous plants to support community livelihoods: the mGU—Useful Plants Project. *J. Environ. Plann. Manage.* <https://doi.org/10.1080/09640568.2016.1166101>.
- Uthpala, T.G.G., Marapana, R.A.U.J., Lakmini, K.P.C., Wettimuny, D.C., 2020. Nutritional Bioactive Compounds and Health Benefits of Fresh and Processed Cucumber (*Cucumis Sativus* L.). *Sumerian J. Biotechnol.* 3 (9), 75–82. <https://doi.org/10.13140/RG.2.2.17510.04161>.
- Uzuozokoro, M.-M.A., Okwesili, F.C.N., Chioma, A.A., 2018. Phytochemical and proximate composition of cucumber (*Cucumis sativus*) fruit from Nsukka, Nigeria. *Afr. J. Biotechnol.* 17 (38), 1215–1219. <https://doi.org/10.5897/ajb2018.16410>.
- van Eck, N.J., Waltman, L., 2010. Software survey: vOSviewer, a computer program for bibliometric mapping. *Scientometrics*. <https://doi.org/10.1007/s11192-009-0146-3>.
- Wilson, D.S., Hartberg, Y., MacDonald, I., Lanman, J.A., Whitehouse, H., 2017. Cultural Diversity Really is Like Biological diversity: Reply to Comments On The Nature of Religious Diversity: A Cultural Ecosystem Approach. *Religion, Brain and Behavior.* <https://doi.org/10.1080/2153599X.2015.1132253>.
- Wink, M., 2013. Evolution of secondary metabolites in legumes (Fabaceae). *S. Afr. J. Bot.* <https://doi.org/10.1016/j.sajb.2013.06.006>.
- Women Jungle Survival Course. (2019).
- Yaseen, G., Ahmad, M., Shinwari, S., Potter, D., Zafar, M., Zhang, G., Shinwari, Z.K., Sultana, S., 2019. Medicinal Plant Diversity Used for Livelihood of Public Health in Deserts and Arid Regions Of Sindh-Pakistan. *Pak. J. Bot.* 51 (2), 657–679.
- Zuhud, E., Sofyan, K., Prasetyo, L., Kartodihardjo, H., 2007. Sikap Masyarakat dan Konservasi: suatu Analisis Kedawung (*Parkia timoriana* (DC) Merr.) sebagai Stimulus Tumbuhan Obat bagi Masyarakat, Kasus di Taman Nasional Meru Betiri (Community's Attitudes and Conservation: an Analysis of Kedawung (*Parkia timoriana* (DC) Merr.) as Stimulus for Medicinal Plants in Meru Betiri National Park). *Jurnal Media Konservasi* XII, 22–32.

Indigenous vegetables consumed as lalapan by Sundanese ethnic group in West Java, Indonesia: Potential, traditions, local knowledge, and it's future

ORIGINALITY REPORT

14%	11%	10%	4%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	journal.unj.ac.id Internet Source	2%
2	"Editorial Board", South African Journal of Botany, 2022 Publication	1%
3	repository.uwc.ac.za Internet Source	1%
4	link.springer.com Internet Source	1%
5	lucris.lub.lu.se Internet Source	<1%
6	Udin Supriadi, Tedi Supriyadi, Aam Abdussalam, Aam Ali. "A Decade of Value Education Model: A Bibliometric Study of Scopus Database in 2011-2020", European Journal of Educational Research, 2022 Publication	<1%
7	www.chacoarchive.org	

<1 %

8

S. DIETZ. "Absciscic acid in lichens: variation, water relations and metabolism", New Phytologist, 1/1998

Publication

<1 %

9

www.scilit.net

Internet Source

<1 %

10

univendspace.univen.ac.za

Internet Source

<1 %

11

N. Salhi, A. Bouyahya, S. Fettach, A. Zellou, Y. Cherrah. "Ethnopharmacological study of medicinal plants used in the treatment of skin burns in occidental Morocco (area of Rabat)", South African Journal of Botany, 2019

Publication

<1 %

12

digitalcommons.unl.edu

Internet Source

<1 %

13

apjtm.org

Internet Source

<1 %

14

worldwidescience.org

Internet Source

<1 %

15

"Ethnobotany of the Mountain Regions of Southeast Asia", Springer Science and Business Media LLC, 2021

Publication

<1 %

16	T.P. Dweba, M.A. Mearns. "Conserving indigenous knowledge as the key to the current and future use of traditional vegetables", International Journal of Information Management, 2011 Publication	<1 %
17	cyberleninka.org Internet Source	<1 %
18	Encyclopaedia of the History of Science Technology and Medicine in Non-Western Cultures, 2016. Publication	<1 %
19	Vita Meylani, Langkah Sembiring, Ahmad Fudholi, Tri Wibawa. "Differentiated sap (4–6) gene expression of Candida albicans isolates from HIV-positive patients with oral candidiasis and commensals in healthy individuals", Microbial Pathogenesis, 2021 Publication	<1 %
20	repository.unisbablitar.ac.id Internet Source	<1 %
21	smujo.id Internet Source	<1 %
22	www.hindawi.com Internet Source	<1 %
23	iiste.org Internet Source	<1 %

24	repository.unam.edu.na Internet Source	<1 %
25	Mohammad Humayoon Amini, Kamran Ashraf, Fatimah Salim, Siong Meng Lim et al. "Important insights from the antimicrobial activity of Calotropis procera", Arabian Journal of Chemistry, 2021 Publication	<1 %
26	egeoscien.neigae.ac.cn Internet Source	<1 %
27	Anne Kathie Samois, Fawzi Mahomoodally. "Ethnopharmacological appraisal of culturally important medicinal plants and polyherbal formulas used against communicable diseases in Rodrigues Island", Journal of Ethnopharmacology, 2016 Publication	<1 %
28	Submitted to Universitas Islam Majapahit Student Paper	<1 %
29	elearning.medistra.ac.id Internet Source	<1 %
30	kobra.uni-kassel.de Internet Source	<1 %
31	stjosephslearningandnews.blogspot.co.nz Internet Source	<1 %

32

Internet Source

<1 %

33

www.jurnal.ar-raniry.ac.id

Internet Source

<1 %

34

Adi Bejo Suwardi, Zidni Ilman Navia.
"Sustainable Use and Management of Wild
Edible Fruit Plants: A Case Study in the Ulu
Masen Protected Forest, West Aceh,
Indonesia", Journal of Sustainable Forestry,
2022

Publication

<1 %

35

es.scribd.com

Internet Source

<1 %

36

ethnobiomed.biomedcentral.com

Internet Source

<1 %

37

www.diva-portal.org

Internet Source

<1 %

38

www.frontiersin.org

Internet Source

<1 %

39

S. S. Ata-Allah. "Cation distribution, hyperfine
parameters and conduction mechanism in the
ferrimagnetic system $\text{Cu}_{0.5}\text{Co}_{0.5}\text{Ga}_x\text{Fe}_{2-x}\text{O}_4$ ",
physica status solidi (b), 05/2005

Publication

<1 %

40

Silvana Piersanti, Valerio Saitta, Manuela
Rebora, Gianandrea Salerno. "Olfaction in

<1 %

phytophagous ladybird beetles: antennal
sensilla and sensitivity to volatiles from host
plants in *Chnootriba elaterii*", *Arthropod-Plant
Interactions*, 2022

Publication

41

repozitorij.hgi-cgs.hr

Internet Source

<1 %

42

www.researchsquare.com

Internet Source

<1 %

43

www.springermedizin.de

Internet Source

<1 %

44

Publication

<1 %

45

"Digital Libraries: Knowledge, Information,
and Data in an Open Access Society", Springer
Science and Business Media LLC, 2016

Publication

<1 %

46

"Ethnobiology of Mountain Communities in
Asia", Springer Science and Business Media
LLC, 2021

Publication

<1 %

47

Bor-Shiuan Cheng. "Paternalistic leadership
and subordinate responses: Establishing a
leadership model in Chinese organizations",
Asian Journal Of Social Psychology, 4/2004

Publication

<1 %

48 Catalina Acuña - Gutiérrez, Víctor M. Jiménez, Joachim Müller. "Occurrence of mycotoxins in pulses", Comprehensive Reviews in Food Science and Food Safety, 2022

Publication

<1 %

49 Mustopa Marli Batubara. "Natural Determinant Stimuli and Agroforestry Benefit Stimuli for Developing Medical Agro-Industry of Kerinci Society in Sebelat Kerinci National Park", Journal of Agricultural Studies, 2017

Publication

<1 %

50 Naomi Bushin. "Quantitative datasets and Children's Geographies: examples and reflections from migration research", Children's Geographies, 2008

Publication

<1 %

51 core-cms.prod.aop.cambridge.org

Internet Source

<1 %

52 journal.frontiersin.org

Internet Source

<1 %

53 scholarworks.waldenu.edu

Internet Source

<1 %

54 www.mdpi.com

Internet Source

<1 %

55 www.tandfonline.com

Internet Source

<1 %

56

A. F. S. L. LOK, K. Y. CHONG, K.-X. TAN, H. T. W. TAN. "A CHECKLIST OF THE SPONTANEOUS EXOTIC VASCULAR PLANT FLORA OF SINGAPORE", COSMOS, 2011

Publication

<1 %

57

Abida Bano, Mushtaq Ahmad, Taibi Ben Hadda, Abdul Saboor et al. "Quantitative ethnomedicinal study of plants used in the skardu valley at high altitude of Karakoram-Himalayan range, Pakistan", Journal of Ethnobiology and Ethnomedicine, 2014

Publication

<1 %

58

journal.pusbindiklatren.bappenas.go.id

Internet Source

<1 %

59

"Indigenous Knowledge and Disaster Risk Reduction", Springer Science and Business Media LLC, 2023

Publication

<1 %

60

"Novel Drug Targets With Traditional Herbal Medicines", Springer Science and Business Media LLC, 2022

Publication

<1 %

61

Raveevatoo Buathong, Sutsawat Duangsrirai. "Plant ingredients in Thai food: a well-rounded diet for natural bioactive associated with medicinal properties", PeerJ, 2023

Publication

<1 %

62

Reza Mareta Putri, Ali Mustadi, Kintan Limiansih. "Analysis of Papuan Students' Learning Motivation in Online Learning for Elementary Science Physics Courses", AL-ISHLAH: Jurnal Pendidikan, 2022

Publication

<1 %

63

Shahana Wahid, Ali Alqahtani, Rafeeq Alam Khan. "Analgesic and anti-inflammatory effects and safety profile of Cucurbita maxima and Cucumis sativus seeds", Saudi Journal of Biological Sciences, 2021

Publication

<1 %

64

eprints.whiterose.ac.uk

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On